No.



9200124

TO ALL TO WHOM THESE PRESENTS SHALL COME;

FFR Cooperative

Colhereus, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLI-CANT(S) FOR THE TERM OF eighteen YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EX-LUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT TY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT . 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

WHEAT

'FFR 555W'

In Testimony Wathercot, I have hereunto set my hand and caused the seal of the Blant Variety Protection Office to be affixed at the City of Washington, D.C.
this 31st day of March in
the year of our Lord one thousand nine
hundred and ninety-three.

Plant Variety Protection Office

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Office, OIRM, Room 404-W, Washington, D.C. 20250; and to the Office of Management and Budget, Paperwork Reduction Project (OMB #0581-0055), Washington, 20250.

U.S. DEPARTMENT OF AGE AGRICULTURAL MARKETIN	RICULTURE IG SERVICE		Application is required in	
APPLICATION FOR PLANT VARIETY (Instructions on re		CERTIFICATE	determine if a plant variety certificate is to be issued (7 U Information is held confide certificate is issued (7 U.S.C. 2	S.C. 2421). Intial until
NAME OF APPLICANT(S) (as it is to appear on the Certificate)		2. TEMPORARY DESIGNATION OR	3. VARIETY NAME	
FFR COOPERATIVE		EXPERIMENTAL NO. VA85-54-290	FFR 555W	vi
4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP)		5. PHONE (Include area code)	FOR OFFICIAL USE O	NLY
			PVPO NUMBER	
P O Box 322			0200124	ł
Battle Ground, IN 47920		(317) 567-2115	9200124	†
·			F Date	
6 CEMIC AND COCCITO MAKE		<u> </u>	March 9 199	<u> </u>
	FAMILY NAME (Botanio	al)	N Time	П
Triticum aestivum	Gramineae		G A.M.	P.M.
8. CROP KIND NAME (Common Name)	9. 1	DATE OF DETERMINATION	F Filing and Examination	Fee:
Wheat, Common		September 1, 1989	S Date	
10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FORM OF ORGANIZ		•	R Dec 31,199	7/
		•	E Certificate Fee:	<u> </u>
Corporation 11. IF INCORPORATED, GIVE STATE OF INCORPORATION	l so De	TE OF INCORPORATION	: 250. ⁵⁰	
	· I		V Date	
Wisconsin		960	5 Mar. 1, 199	73
13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO SE	ERVE IN THIS APPLICATION	N AND RECEIVE ALL PAPERS		
Frank Spelbring FFR COOPERATIVE				
P O BOX 322		(317) 567-21	15	
Battle Ground, IN 47920		· ·		
14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow	INSTRUCTIONS on rever	PHONE (Include area cod	<i>ae)</i> :	
a. X Exhibit A, Origin and Breeding History of the Variety.		•		
b. X Exhibit B, Novelty Statement.				
c. X Exhibit C, Objective Description of Variety.				
d. X Exhibit D, Additional Description of Variety.				7. T.
e. X Exhibit E, Statement of the Basis of Applicant's Ownership.		•		
1. X Seed Sample (2,500 viable untreated seeds). Date Seed Se			 •	5.5
g. X Filing and Examination Fee (\$2,150) made payable to "Tre. 15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VARIETY BE SOLD			and an extension of the Division of the state of the stat	
Protection Act.) YES (If "YES." answer items 16 and 17 below	-	AS A CLASS OF CERTIFIED SEED? (SE D," skip to item 18 below)	ee section 63(a) of the Plant Variet	y ;
16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS?		TEM 16, WHICH CLASSES OF PRODU	JCTION BEYOND BREEDER SEED?	,
		_		
YES NO	i ∐ Fo∪	NDATION REGIST	TERED CERT	TIFIED
18. DID THE APPLICANT(S) PREVIOUSLY FILE FOR PROTECTION OF THE VARIE	TY IN THE U.S.?			
YES (If "YES," through Plant Variety Protection Act NO	Patent Act. Give dat	e:		
19. HAS THE VARIETY BEEN RELEASED, USED, OFFERED FOR SALE, OR MAF	OVETED IN THE U.S. CO.	THER COUNTRIES		
		BER 20, 1991		SE S
				19
20. The applicant(s) declare(s) that a viable sample of basic seed request in accordance with such regulations as may be applications.	ıble.		-	
The undersigned applicant(s) is (are) the owner(s) of this se uniform, and stable as required in section 41, and is entitled t	xually reproduced no protection under th	ovel plant variety, and believe	e(s) that the variety is dist Plant Variety Protection Ac	.inct,
Applicant(s) is (are) informed that false representation herein	i can jeopardize prote	ction and result in penalties.		
SIGNATURE OF APPLICANT JOURNALS)	CAPACITY OR T		DATE	
	Mas 1			
Trank Spellering	Mark	ting Deserto	1/2-20-9	7 /
SIGNATURE OF APPLICANT (Owner(s))	CAPACITY OR T	TLE /	DATE	

Wheat 'FFR 555W'

14A. Exhibit A: Parentage

FFR 555W was derived from one of six crosses in which 'Coker 76-35' was a common parent. F_1 seed of the six crosses including Coker 76-35/'Doublecrop', Coker 76-35/Va. 72-54-14, Coker76-35/Va. 76-52-12, Coker 76-35/Va. 76-52-24, Coker 76-35//Coker 65-20/Arthur, and Coker 76-35/3/(Kavkaz//Coker 65-20/Arthur), F₁ was composited and advanced as a bulk population.

Va. 72-54-14 is a selection from the cross 'Blueboy'/3/'Atlas 50'/1950 Row 223//'Redcoat' or Redcoat sib. The 1950 Row 223 parent had leaf and/or stem rust resistance. Va. 76-52-12 is a selection from the cross 'Blueboy'/'Knox 62'. Va. 76-52-24 is a selection from the cross 'Blueboy'//Thorne $^*5/199-4/3/Blueboy$ Sel. The 199-4 parent was the F, from the cross of 'Asosan'/3/'Supresa'/'Redhart'//'Chancellor' with P55-47.1-5, which was a line in which the leaf rust resistance of Aegilops umbellulata was transferred into 'Chinese Spring'.

FFR 555W was selected in 1984 as a F_{θ} headrow, using a modified bulk breeding system. This selection was grown in an observation plot in 1985 as entry 290 in test 54. This line, designated as Va. 85-54-290, has been evaluated in the Virginia State Variety Trials since 1987.

A large increase block of Va. 85-54-290, approximately 64 ft. by 66 ft., was planted in 1987, rogued thoroughly for aberrant types, and harvested in 1988. Seed from this block was planted at the Foundation Seed Farm in 1989, and rogued to remove gross off-types. current lot of Foundation seed (F_{12} generation), derived from this multiplication, still contains a small percentage of aberrant types, primarily later maturing, taller plants with dense heads, blue-colored glumes, and long rachis hairs. These off-type plants do not exceed 5% Voucout of the population, are readily identifiable, and are within acceptable limits for seed certification. FFR 555W is genetically stable in the sense that the variety can be maintained and reproduced via seed sense that the variety can be maintained and reproduced via seed without changing its characteristics.

Approximately 276 heads were selected from the 1987-88 increase block for use in establishing an improved lot of Breeder seed. These heads were threshed individually, and grown as headrows in 1988-89. Of the 276 headrows, 238 were saved and planted in individual six-row plots, three feet in length. A sample of seed from each headrow was also used to test each row for seedling reaction to a mixture of two mildew cultures, and to a single race of leaf rust. Of the 238 headrows evaluated, 227 had an intermediate to moderately susceptible reaction to mildew, and were susceptible to leaf rust. Upon consideration of greenhouse and field evaluations, 233 of the 238 plots were harvested and bulked. This Breeder seed of FFR 555W was provided to the Foundation Seed Farm, and will be the source of future seed multiplications. Within the limits of biological expectation, the Breeder seed of FFR 555W is uniform and stable.

FFR 555W Wheat

14B. Exhibit B: Novelty Statement

FFR 555W most closely resembles 'Coker 983', which is a sister line of Coker 76-35. FFR 555W has a red coleoptile color, while that of Coker 983 is white. Based on data from approximately 20 tests conducted over four years, FFR 555W was four inches taller and had a 2.6 lb/bu lower test weight than Coker 983. FFR 555W and Coker 983 express a similar level of resistance to powdery mildew (Erysiphe graminis DC f. sp. tritici E. Marchal) in the field; however, FFR 555W has a higher level of seedling resistance than Coker 983 based on their differential reaction to two mildew cultures with different virulence formulas. FFR 555W is moderately resistant to a mildew culture which is virulent on Pm4a and Pm7, and avirulent on Pm1, Pm2, Pm3a, Pm3b, Pm3c, Pm2 + Pm6, Pm8, and PmMA, while Coker 983 is moderately susceptible to this culture. FFR 555W also expresses an intermediate to moderate level of resistance to a mildew culture which is virulent on Pm2, Pm3a, Pm3c, Pm4a, and PmMA, and avirulent on Pm1 and Pm3b, while Coker 983 is susceptible to this culture. Evaluations at the Cereal Rust Laboratory, St. Paul, MN, indicate that FFR 555W has the Lr10 gene for leaf rust (<u>Puccinia recondita</u> Rob. ex Desm. f. sp. <u>tritici</u>) resistance, while Coker 983 has <u>Lr18</u> in addition to <u>Lr10</u>. Further evaluations at the Cereal Rust Laboratory, have shown that FFR 555W is resistant to the stem rust (Puccinia graminis Pers. f. sp. tritici) race RKQS which is virulent on Sr5, Sr9b, Sr7b, Sr6, Sr8a, Sr9a, Sr36, Sr9b, Sr15, Sr16, and Sr13 + <u>Sr17</u>, and avirulent on <u>Sr9e, Sr11, Sr13, Sr10</u>, and <u>SrTMP</u>, while Coker 983 is susceptible to this race.

EXHIBIT C

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK AND SEED DIVISION BELTSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY

INSTRUCTIONS: See Reverse. WHEAT (TR	FOR OFFICIAL USE ONLY
FFR COOPERATIVE	SVSO NUMBER
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	9200124
4112 E. State Road 225	VARIETY NAME OR YEMPORARY DESIGNATION
W. Lafayette, IN 47906	DESIGNATION
	FFR 555W
Place the appropriate number that describes the varietal character Place a zero in first box (e-s- 0 8 9 or 0 9) when number	r of this variety in the boxes below. is either 99 or less or 9 or less.
I. KIND:	
1 1 = COMMON 2 = DURUM 3 = EMMER 4 = SPELT 5	= POLISH 6 = POULARD 7 = CLUB
2. TYPE,	1 = SOFT 3 = OTHER (Specify)
2 1 = SPRING 2 = WINTER 3 = OTHER (Specify)	1 2 = HARD
2 1 = WHITE 2 = RED 3 = OTHER (Specify)	
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	
FIRST FLOWERING	LAST FLOWERING
4. MATURITY (50% Flowering):	
0 2 NO. OF DAYS EARLIER THAN	7 1 = ARTHUR 2 = SCOUT 3 = CHRIS
0 5 NO. OF DAYS LATER THAN	8 4 = LEMHI 5 = NUGAINES 6 = LEEDS 7 = TYLER 8 = COKER 916
5. PLANT HEIGHT (From soil level to top of head):	, Trees, o ones
0 9 8 cm. HIGH	
0 6 CM. TALLER THAN	8 7 = TYLER 8 = COKER 916
0 9 CM. SHORTER THAN	7 1 = ARTHUR 2 = SCOUT 3 = CHILDS 4 = LEMHI 5 = NUGAINES 6 = LEEDS
6. PLANT COLOR AT BOOTING (See reverse):	7. ANTHER COLOR: Red at base of stamen
3 1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN	1 1 = YELLOW 2 = PURPLE
8. STEM:	
Anthocyanin: I = ABSENT 2 = PRESENT	2 Waxy bloom: I = ABSENT 2 = PRESENT
Hairiness of last No hairs on Peduncle internode of rachis: 1 = ABSENT 2 = PRESENT	1 Internodes: 1 = HOLLOW 2 = SOLID
0 3 (sometimes 4) (on margins)	2 2 CM. INTERNODE LENGTH BETWEEN FLAG LEAF
9. AURICLES:	
Anthocyanin: 1 = ABSENT 2 = PRESENT	2 Hairiness: 1 = ABSENT 2 = PRESENT
O. LEAF:	
Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):	1 Flag leaf: 1 = NOT TWISTED 2 = TWISTED
Hairs of first leaf sheath: = ABSENT 2 = PRESENT	2 Wazy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT
1 2 MM. LEAF WIDTH (First leaf below flag leaf)	2 1 CM. LEAF LENGTH (First leaf below flag leaf):

2 3 = MIDE (CA. 9 mm.) 8.2 - 8.7 mm 2 3 = MIDE (CA. 4 mm.) 3.6 - 3.8 mm				
1 = WHITE	. -			
2 Color at maturity 5; BROWN 6 = BLACK 7 = OTHER (Specify: 1 1 1 MM. WIOTH 12 CLUMES AT MATURITY: 2 LEARS 1 = SHORT (CA. 7 mm.) 2 = MEDIUM (CA. 2 mm.) 3 = CONG (CA. 9 mm.) 8.2 - 8.7 mm 2 Leads 1 = SHORT (CA. 7 mm.) 2 = MEDIUM (CA. 3 mm.) 3 = WIOE (CA. 9 mm.) 3.5 - 3.8 mm 3 = CONG (CA. 9 mm.) 2 = MOUNDED 2 Shoulder 1 = WANTING 7 = OBLIQUE 3 = MOUNDED 2 Shoulder 1 = WANTING 7 = OBLIQUE 3 = MOUNDED 2 HAVE 1 = CONG (CA. 9 mm.) 2 = MEDIUM (CA. 3.3.3		3 Awnedness: 1 = AWNLESS 2 = APICALLY AWNLETED	3 = AWNLETED 4 = AWNED	·
12. GLUMES AT MATURITY: Legath:		2 Color at maturity: 5 = BROWN 6 = BLACK 7 = OTH		·
2 Length:		0 9 CM. LENGTH	1 1 мм. wютн	
2- 4 Shoulder 1= WANTING 2= OBLIQUE 3 = ROUNDED 1 Serie: 1= OBTUSE 2 = ACUTE 3 = ACUMINATE 13. COLEOPTILE COLOR: 2 1 = WHITE 2 = RED 3 = PURPLE		Length: 1 = SHORT (CA. 7 mm.) 2 = MEDIUM (CA. 8 mm.)	1 01	2 6 2 0
1 = ABSENT 2 = PRESENT 2 = PRESENT 15. JUVENILE PLANT GROWTH HABIT: 2 1 = PROSTRATE 2 = SEMI-ERECT 3 = ERECT 16. SEED: 1 Shape: 1 = OVATE 2 = OVAL 3 = ELLIPTICAL 1 Check: 1 = ROUNDED 2 = ANGULAR W/ flange of endosperm on cheel 2 Brush: 1 = NOT COLLARED 2 = COLLARED 2 Brush: 1 = NOT COLLARED 2 = COLLARED 3 = LONG 2 Brush: 1 = NOT COLLARED 2 = COLLARED 3 = COL	2-	Shoulder 1 = WANTING 2 = OBLIQUE 3 = ROUNDED	Beak: 1 = OBTUSE	2 = ACUTE 3 = ACUMINATE
15. JUVENILE PLANT GROWTH HABIT: 2 1 = PROSTRATE 2 = SEMI-ERECT 3 = ERECT 16. SEED: 1 Shape: 1 = OVATE 2 = OVAL 3 = ELLIPTICAL 1 Cheek: 1 = ROUNDED 2 = ANGULAR W/ flange of endosperm on cheel 2 Brush: 1 = NOT COLLARED 2 = COLLARED 2 = COLLARED 3 = LONG 2 Brush: 1 = NOT COLLARED 2 = COLLARED 3 = LONG 2 Brush: 1 = NOT COLLARED 2 = COLLARED 3 = LONG 3 3 3 3 3 3 3 3 3		13. COLEOPTILE COLOR:	14. SEEDLING ANTHOCYA	NIN:
2 1 = PROSTRATE 2 = SEMI-ERECT 3 = ERECT 3 = ERECT 16. SEED: 1	٠	2 1 = WHITE 2 = RED 3 = PURPLE	1 = ABSENT 2 =	PRESENT
16. SEED: 1 Shape: 1=OVATE 2=OVAL 3=ELLIPTICAL 2 Brush: 1=SHORT 2=MEDIUM 3=LONG 2 Brush: 1=SHORT 2=MEDIUM 3=LONG 4 Phenol reaction 1=IVORY 2=FAWN 3=LT. BROWN 4 = BROWN 5=BLACK 3 Colur: 1=WHITE 2=AMBER 3=RED 4=PURPLE 5=OTHER (Specify) 0 6 MM. LENGTH 0 3 MM. WIDTH 3 9 GM. PER 1000 SEEDS 17. SEED CREASE: Width: 1= 603 OR LESS OF KERNEL 'WINOKA' 18. DISEASE: (0 = Not Tested, 1 = Susceptible, 2 = Resiston) 19. INSECT: (0 = Not Tested, 1 = Susceptible, 2 = Resiston) Audit-Plant Resistante 19. INSECT: (0 = Not Tested, 1 = Susceptible, 2 = Resiston) O SAWFLY O APHID (Brydw.) 10 O GREEN BUG O CEREAL LEAF BEETLE CMARACTER NAME OF VARIETY NAME OF VARIETY CHARACTER NAME OF VARIETY NAME OF VARIETY CHARACTER NAME OF VARIETY CLeaf corriage Seed shape Leaf corriage Seed shape Check: 1 = ROUNDEO 2 = ANGULAR W/ flange of endosperm on cheel 2 Brush: 1 = NOT COLLARED Slightly A colurity is not collared 2 = COLLARED Slightly Adultable 2 = COLLARED Slightly Seed shape 1 OTHER (Specify) Adultable 1 OTHER (Specify) RACE: L Seed shape 1 OTHER (Specify) RACE: C CHARACTER NAME OF VARIETY CHARACTER CHARACTER		15. JUVENILE PLANT GROWTH HABIT:		
1 Shape: i = OVATE 2 = OVAL 3 = ELLIPTICAL 2 Brush: l = SHORT 2 = MEDIUM 3 = LONG 4 Phenol reaction 1 = IVORY 2 = FAWN 3 = LT. BROWN 4 = BROWN 5 = BLACK 3 Color: l = WHITE 2 = AMBER 3 = RED 4 = PURPLE 5 = OTHER (Specily) 0 6 MM. LENGTH 0 3 MM. WIDTH 3 9 GAR. PER 1000 SEEDS 17. SEED CREASE: Width: l = 802 OR LESS OF KERNEL 'WINOKA' 2 = 802 OR LESS OF KERNEL 'CHRIS' Narrow 2 = 802 OR LESS OF KERNEL 'LEMHI' 18. DISEASE: (0 = Not Tested, l = Susceptible, 2 = Resistant) 2 = STEN RUST (Races) Broadly 1 LEAF RUST Lr 10* 0 STRIPE RUST (Races) Broadly 1 LeOSE SMUT 2 = OVERANT ON APHID (Bydn.) 19. INSECT: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) OTHER (Specily) Hessian Fly Race: L RACE: O D D D D D D D D D D D D D D D D D D		2 1 = PROSTRATE 2 = SEMI-ERECT 3 = ERE	СТ	
W/ flange of endosperm on cheel 2 Brush. = SHORT 2 = MEDIUM 3 = LONG 2 Brush. 1 = NOT COLLARED 2 = COLLARED		16. SEED:	·	
2 Brush. 1 = SHORT 2 = MEDIUM 3 = LONG 4 Phenol reaction 1 = (VORY 2 = FAWN 3 = LT. BROWN 5 = BLACK 3 Color: 1 = WHITE 2 = AMBER 3 = RED 4 = PURPLE 5 = OTHER (Specify) 0 6 MM. LENGTH 0 3 MM. WIDTH 3 9 GM. PER 1000 SEEDS 17. SEED CREASE: Width: 1 = 80% OR LESS OF KERNEL 'WINDKA' 18. DISEASE: (0 = Not Tasted, 1 = Susceptible, 2 = Resistann) *Genes proposed for said Variety by Cereal Ru: 2 **STEM RUST** Resistant** 1 **DISEASE: (0 = Not Tasted, 1 = Susceptible, 2 = Resistann) *Genes proposed for said Variety by Cereal Ru: 2 **Resistant** 1 **OTHER (Specify) **Adult-Plant Resistance** 1 **OTHER (Specify) **HESSIAN FLY RACES: (0 = Not Tasted, 1 = Susceptible, 2 = Resistann) **OTHER (Specify) **Adult-Plant Resistance** 1 **OTHER (Specify) **HESSIAN FLY RACES: (0 = Not Tasted, 1 = Susceptible, 2 = Resistann) **OTHER (Specify) **Adult-Plant Resistance** 1 **OTHER (Specify) **HESSIAN FLY RACES: (0 = Not Tasted, 1 = Susceptible, 2 = Resistann) **OTHER (Specify) **Adult-Plant Resistance** 1 **OTHER (Specify) **HESSIAN FLY RACES: (0 = Not Tasted, 1 = Susceptible, 2 = Resistann) **OTHER (Specify) **OTHER (S		1 Shape: 1 = OVATE 2 = OVAL 3 = ELLIPTICAL	7 - 1	
A Phenoi reaction 1 = IVORY 2 = FAWN 3 = LT. BROWN 5 = BLACK 3 Colur: 1 = WHITE 2 = AMBER 3 = RED 4 = PURPLE 5 = OTHER (Specify)		2 Brush. 1 = SHORT 2 = MEDIUM 3 = LONG		LARED 2 = COLLARED
17. SEED CREASE: Width: = 60% OR LESS OF KERNEL 'WINOKA' Depth: = 20% OR LESS OF KERNEL 'SCOUT'		-1 A 1	IN .	Silghtly
17. SEED CREASE: Width: = 60% OR LESS OF KERNEL 'WINOKA' NATTOW 2 = 80% OR LESS OF KERNEL 'CHRIS' NATTOW 3 = NEARLY AS WIDE' AS KERNEL 'LEMHI' 18. DISEASE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) *Genes proposed for Said Variety by Cereal Rules, (Races) Broadly 1 (Races) Lr 10* 0 STRIPE RUST 0 Loose SMUT 0 LOOSE SMUT 0 SAWFLY 0 BUNT 0 GREEN BUG 0 CEREAL LEAF BEETLE 1 OTHER (Specify) Hessian Fly Race: L RACES: 0 D D D D D D D D D D D D D D D D D D		3 Color: 1 = WHITE 2 = AMBER 3 = RED 4 = PURPLE	5 = OTHER (Specity)	
17. SEED CREASE: Width: = 60% OR LESS OF KERNEL 'WINOKA' 2 = 30% OR LESS OF KERNEL 'SCOUT' Narrow 2 = 80% OR LESS OF KERNEL 'CHRIS' Middeep 3 = 50% OR LESS OF KERNEL 'CHRIS' 18. DISEASE: (0 = Not Tested, 1 = Susceptible, 2 = Resistont) *Genes proposed for said Variety by Cereal Rul 2 STEM RUST			3 9 GM. PER 1000 S	FEDS
Width: 1 = 50% OR LESS OF KERNEL 'WINOKA' Narrow 2 = 30% OR LESS OF KERNEL 'CHRIS' Narrow 3 = NEARLY AS WIDE' AS KERNEL 'LEMHI' 18. DISEASE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) Provided Provide		O O MM. CENGIA	<u> </u>	
Narrow 2 = 80% or Less of Kernel 'Chris' Narrow 3 = NEARLY AS WIDE'AS KERNEL 'LEMHI' 18. DISEASE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) *Genes proposed for said Variety by Cereal Rust (Racco) Broadly 1 (Racco) Lab, St. Paul, (Racco) Broadly 1 (Racco) Lab, St. Paul, (Racco) Broadly 1 (Racco) United (United (United (Racco) United (United (Unit				
Narrow 3 = NEARLY AS WIDE AS KERNEL 'LEMH!' 18. DISEASE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) *Genes proposed for said Variety by Cereal Rust (Racce) Broadly 1 LEAF RUST LAD, St. Paul, Resistant* D BUNT D CREEN BUG D COOSE SMUT 2 POWDERY MILDEW Adult-Plant Resistance 19. INSECT: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) D GREEN BUG D CEREAL LEAF BEETLE 1 OTHER (Specify) Hessian Fly Race: L Racces: D D D D D 20. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER NAME OF VARIETY CHARACTER NAME OF VARIETY CHARACTER NAME OF VARIETY CHARACTER NAME OF VARIETY Celegical Color Coleoptile elongation Leaf carriage Seed shape Celegical Color Coleoptile elongation Seedling pigmentation		1 1	1	
18. DISEASE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) 2	N		Middoon	
2 STEM RUST (Races) Broadly 1 (Races) Ly 10* 0 STRIPE RUST (Races) Broadly 1 (Races) Ly 10* 0 LOOSE SMUT 2 POWDERY MILDEW DEW Adult-Plant Resistance 19. INSECT: (0 = Nor Tested, 1 = Susceptible, 2 = Resistant) 0 SAWFLY 0 APHID (Bydv.) 0 GREEN BUG 0 CEREAL LEAF BEETLE 1 OTHER (Specify) Hessian Fly Race: L RACES: 0 D 1 E 0 F 0 G 20. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER NAME OF VARIETY CHARACTER NAME OF VARIETY Plant tillering Seed size Leaf size Seed shape Leaf color Coleoptife elongation Leaf carriage Seedling pigmentation		J = NEARLY AS WIDE AS KERNEL "EMO!		
2 (Reces) Broadly 1 (Reces) Lr 10* 0 (Reces) Resistant* 0 BUNT OTHER (Specify) Adult-Plant Resistance 19. INSECT: (0 = Nor Tested, 1 = Susceptible, 2 = Resistant) O SAWFLY 0 APHID (Bydn.) 0 GREEN BUG 0 CEREAL LEAF BEETLE 1 OTHER (Specify) Hessian Fly Race: L RACES: 0 D D 1 E 0 F 0 G 20. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER NAME OF VARIETY CHARACTER NAME OF VARIETY Plant tillering Seed size Leaf size Seed shape Leaf color Coleoptile elongation Leaf corriage Seedling pigmentation	:			Lab. St. Paul, MN
Resistant* O BUNT Adult-Plant Resistance 19. INSECT: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) O SAWFLY O APHID (Bydw.) O O GREEN BUG O CEREAL LEAF BEETLE TOTHER (Specify) Hessian Fly Race: L RACES: O D T O D O GREEN BUG O CEREAL LEAF BEETLE T O D O D O D O D O D O D O D	٠.	2 (Races) Broadly 1 (Races) Lr 10*		U LOOSE SMUT
19. INSECT: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) 0 SAWFLY 0 APHID (Bydw.) 1 OTHER (Specify) Hessian Fly Race: L 1 OTHER (Specify) Hessian Fly Race: L 1 OF O C 20. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER NAME OF VARIETY CHARACTER NAME OF VARIETY Plant tillering Seed size Leaf size Seed shape Leaf color Coleoptile elongation Leaf carriage Seedling pigmentation		Resistant* O BUNT	OTHER (Specify)	
O SAWFLY O APHID (Bydv.) O GREEN BUG O CEREAL LEAF BEETLE O THER (Specify) Hessian Fly Race: L RACE: D O D O D O D O D O D O D O D				
1 OTHER (Specity) Hessian Fly Race: Leaf size Leaf carriage Hessian Fly Resembles that Submitted: Character Name of Variety Character Name of Variety Caleoptile elongation Seed ling pigmentation	•	<u></u>	[O]	0 000041 1515 055715
Race: L RACES: O D 1 E O F O G 20. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER NAME OF VARIETY CHARACTER NAME OF VARIETY Plant tillering Seed size Leaf size Seed shape Leaf color Coleoptife elongation Leaf corriage Seedling pigmentation		O SAWFLY U APHID (Bydv.)	GREEN BUG	O CEREAL CEAR BEETER
RACE: RACES: ODD 1 E OF OG 20. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER NAME OF VARIETY CHARACTER NAME OF VARIETY Plant tillering Seed size Leaf size Seed shape Leaf color Coleoptile elongation Leaf corriage Seedling pigmentation		1 OTHER (Specify) Hessian Fly HESSIAN FLY		1 в Ос
20. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THAT SUBMITTED: CHARACTER NAME OF VARIETY CHARACTER NAME OF VARIETY Plant tillering Seed size Leaf size Seed shape Leaf color Coleoptite elongation Leaf corriage Seedling pigmentation		Care: I		
CHARACTER NAME OF VARIETY Plant tillering Leaf size Leaf color Leaf carriage CHARACTER NAME OF VARIETY Seed size Coleoptile elongation Seedling pigmentation			0 0 1 =	0 F 0 G
Plant tillering Seed size Leaf size Seed shape Leaf color Coleoptite elongation Leaf carriage Seedling pigmentation		20. INDICATE WHICH VARIETY MOST CLOSELY RESEMBLES THAT	· y ·································· ···	
Leaf size Seed shape Leaf color Caleoptile elongation Leaf carriage Seedling pigmentation		The second secon	{	NAME OF VARIETY
Leaf color Coleoptile elongation Leaf carriage Seedling pigmentation			<u> </u>	
Leaf carriage Seedling pigmentation		The state of the s		
	,		 	
INCTDICTIONS	•		<u> </u>	

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

- (a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Wheat Varieties Grown in the United States, Technical Bulletin 1278, United States Department of Agriculture.
- (b) W.E. Walls, 1965. A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity. contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)

FFR 555W Wheat

14D. Exhibit D: Additional Description of FFR 555W

Since FFR 555W has not been tested in comparison with any of the six cultivars indicated for wheat in Exhibit C, data on its performance in Virginia over a period of four years (1987-1990) are presented in the tables which follow this section.

FFR 555W was also evaluated in the 1989-90 Uniform Southern Soft Red Winter Wheat Nursery. Performance data are summarized in the USDA nursery report compiled by Dr. Harold Bockelman.

Quality evaluations made at the Soft Wheat Quality Laboratory, Wooster, OH, indicate that FFR 555W has good milling and baking properties compared with standard checks. Quality data for FFR 555W are presented in the tables which follow this section.

· .			G	RAIN Y bu/ac			
	1990 (6)	1989 (6)	1988 (8)	1987 (6)	1986 (7)	1987-90 (26)	1986-90 (33)
Madison	71	73	87	65	54	74	70
Wakefield	69	- 80	93	67	58	77	73
FFR 555W	84	79	91	68		81	
Massey	71	73	77	63	51	. 71	67
Tyler	69	64	71	53	52	64	62
Saluda	70	66	<i>7</i> 1	57	50	66	63
Florida 302	76	76	82	64	53	7 5	70
Coker 916	67	65	82	62	50	69	65
Coker 983	<i>7</i> 5	71	84	66	51	74	69
Coker 833	<i>7</i> 5	64	<i>7</i> 5	62	51	69	65
Pioneer 2550	70	64	78	53	49	66	63
Pioneer 2555	68	68	84	60		7 0	
Pioneer 2548	79	69			****		
FFR 568	· 75	69					

8.6

6.5

L.S.D. (0.05)

4.0

	. 4.		TES	T WE!			
	1990 (4)	1989 (6)	1988 (8)	1987 (6)	1986 (7)	1987-90 (24)	1986-90 (31)
Madison	58.1	55.9	58.0	55.4	59.6	56.9	57.4
Wakefield	57.7	55.7	58.3	55.3	59.6	56.8	<i>57.</i> 3
FFR 555W	57.9	55.7	58.3	53.8		56.4	
Massey	59.0	57.4	58.9	57.0	60.1	58.1	58.5
Tyler	57.0	55.5	57.9	54.4	58.8	56.2	56.7
Saluda	60.7	57.2	60.2	56.7	62.2	58.7	59.4
Florida 302	57.2	54.2	57.9	54.3	59.1	55.9	56.5
Coker 916	55.8	55.4	58.5	55.6	60.3	56.3	57.1
Coker 983	59.9	57.8	60.2	57.9	61.3	59.0	59.4
Coker 833	58.0	55.6	58.2	56.0	58.6	57.0	57.3
Pioneer 2550	57.5	55.1	58.9	55.0	60.1	56.6	57.3
Pioneer 2555	57.7	54.8	58.2	54.5	*****	56.3	
Pioneer 2548	58.1	54.2		****			
FFR 568	58.3	56.0					
L.S.D. (0.05)			1.0	1.2	0.7		

[†]The number in parentheses below column headings indicates the number of tests on which data are based.

Table 1a. Average Performance of Wheat Cultivars Evaluated in Virginia, 1986-90.†

9200124

DATE HEADED	
MAR 31+	

	1990 (3)	1989 (3)	1988 (6)	198 7 (5)	1986 (5)	1987-90 (17)	1986-90 (22)
Madison	26	35	34	40	29	34	33
Wakefield	32	38	39	43	35	38	37
FFR 555W	28	38	40	42		37	
Massey	29	37	37	41	33	36	35
Tyler	32	40	41	44	36	39	39
Saluda	28	38	38	41	33	36	36
Florida 302	30	37	38	42	35	37	36
Coker 916	25	33	34	38	28	33	32
Coker 983	29	37	37	42	33	36	36
Coker 833	33	41	42	44	36	40	39
Pioneer 2550	32	39	41	42	36	39	38
Pioneer 2555	28	37	37	40		36	
Pioneer 2548	30	37					
FFR 568	31	38			***************************************		
L.S.D. (0.05)			2.0	1.3	2.0		

PLANT HEIGHT Inches

	1990 (3)	1989 (3)	1988 (8)	1987 (6)	1986 (7)	1987-90 (20)	1986-90 (27)
Madison	36	39	41	41	34	39	38
Wakefield	38	41	43	42	35	41	40
FFR 555W	36	. 38	41	40		39	
Massey	38	41	43	43	36	41	40
Tyler	39	43	45	43	37	43	41
Saluda	34	36	39	38	31	37	36
Florida 302	39	41	42	41	33	41	39
Coker 916	34	36	39	37	32	37	36
Coker 983	33	35	36	36	29	.35	34
Coker 833	38	41	43	42	34	41	40
Pioneer 2550	37	39	42	40	32	40	38
Pioneer 2555	36	38	37	40		38	****
Pioneer 2548	35	36			****	****	
FFR 568	39	40	·				
L.S.D. (0.05)			1.4	1.0	2.0		

[†]The number in parentheses below column headings indicates the number of tests on which data are based.

Table 1b. Average Performance of Wheat Cultivars Evaluated in Virginia, 1986-90.†

			LOI	OGING %			Winter Surviva %
	1990 (2)	1989 (5)	1988 (7)	1987 (4)	1986 (3)	1987-90 (18)	1988 (1)
Madison	13	9	14	35	0	18	100
Wakefield	14	16	21	29	0	20	100
FFR 555W	8	9	8	25		13	100
Massey	30	23	27	46	1	32	100
Tyler	16	10	11	25	0	16	100
Saluda	11	21	29	43	0	26	100
Florida 302	6	5	5	35	0	13	25
Coker 916	20	24	19	32	3	24	100
Coker 983	8	10	3	24	0	- 11	73
Coker 833	21	36	11	31	3	25	100
Pioneer 2550	21	15	22	25	0	21	100
Pioneer 2555	7	6	4	15		8	100
Pioneer 2548	6	5					
FFR 568	7	7		· ·			
L.S.D. (0.05)		·	14	22			

[†]The number in parentheses below column headings indicates the number of tests on which data are based.

Table 2. Reaction of Wheat Cultivars to Diseases in Virginia, 1986-90.+

		Powd	Powdery Mildew	ildew			Leaf Rust	Rust		So	Soilborne Viruses	e Virus	es
			%				%			0-5‡	Ì	%	1
	1990 (5)	1989	1988 (7)	1987	1986 (5)	1990 (5)	1988 (2)	1987 (1)	1986 (1)	1990 (1)	1988	1987	1986
Madison	16	9	r-i	1	-80	8	1	-	17	0	1		-
Wakefield	46	7	0	0	Ŋ	9	က	22	7	က	09	ß	
FFR 555W		က	∞	4	1	26	14	15		ю	40	₩	
Massey	6	4	4	် က		40	39	38	33	0	-	0	0
Tyler	42	25	32	32	15	37	45	22	47	ဗ	10	10	H
Saluda	56	39	40	40	38	11	15	8	20	ī.	100	- 8	70
Florida 302	17	4	S	1		4	0	0	-	4	100	08 ·	86
Coker 916	12	∞	6	7	2	9	Н	4	0	က	100	7	17
Coker 983	7	2	ဇ	Т	, -	38	က	0	4	რ	100	17	H
Coker 833	17	12	18	13	∞	0	ເດ	5	0	₩	2	₩	2
Pioneer 2550	29	14	30	25	20	26	11	9	Η .	4	63	13	63
Pioneer 2555	%	18	28	24		4	0	7	-	4	0	0	
Pioneer 2548	24	10	1 1		-	₩		8 k 1	!	5		l	1
FFR 568	14	Ŋ	İ	1	I	7	1		1	H	1		
L.S.D. (0.05)		1	6	10	6		13		ŀ				

+ The number in parentheses below column headings indicates the number of tests on which data are based.

0=Resistant; 5=Susceptible

Table 3. Soft Wheat Milling and Baking Quality Evaluations for 'FFR 555W' Wheat, 1987-1989+.

,	Milling Qual.		Baking Qual.		Break	St. Gr.	Straigh Fle	Straight Grade Flour	Micro	Cookie						
Entry	Score/ Grade		ore/ Grade	Test Wt.	Flour Yield	Flour	Flour Ash %	Flour Pro. %	AWRC	Diameter CM.	Top Grain	Soft. Equiv.	Red	Fria- bility	EST.	Mill- ability
1987 Virginia																
FFR 555W	99.1 I	В 93.1	.1 C	74.1		72.2	.35	6.9	51.7	18.0	9	64.6				
Saluda (Std.)	100 A	A 100	0 A	77.8		70.4	.31	6.9	54.3	18.4	9	68.4		-		
Tyler	101.8 ₽	A 98.7	.7 B	73.3		71.3	-3	6.7	53	18.1	9	2.29				
Massey	101.6 A	A 94.6	.6 C	76.4		71.7	.33	7.5	53.5	18.2	9	65.4				
LSD				1.17		0.8	0.035	0.56	1.49	0.25	2.06	3.7				
1988 Virginia						:										
FFR 555W	100. ⊿	A 11	110.2 A	62	26.4	78.1	.42	9.05	51.6	17.86	S		7	29.2	9.1	114.7
Saluda (Std.)	100	A 100	0 A	63.2	29.1	76.4	.37	6.7	54.4	17.73	2		7	27.9	10.3	111.1
Tyler	97.1 I	B 91.9	.9 C	61.8	27.3	76.1	36	9.34	52.1	17.44	4		7	27.6	11.2	108.7
LSD				0.58	0.748	0.90	0.02	0.54	1.74	0.24	0.83			0.33	0.95	9.22
1989 Virginia																
FFR 555W	101.7	A 10	106.0 A	74.7		76.3		7.70	53.7	18.2	9	59.1				
Massey (Std.)	100.0	A 10	100.0 A	7.92		75.4		8.87	55.6	17.9	4	59.8				
Tyler	99.1	B 10	106.9 A	75.0		75.3		7.61	55.3	18.05	5	60.7				
LSD				1.155		0.81		0.718	1.529	0.239		3.216				

+ Milling and baking evaluations performed by the USDA Soft Wheat Quality Laboratory at Wooster, OH.

Exhibit E. Statement of the Basis of Applicant's Ownership

FFR 555W was bred and developed by plant Breeders at Virginia Polytechnic Insititute and State University (VPI) at Blacksburg, Virginia. Under an agreement between FFR cooperative and VPI, exclusive, proprietary rights were granted to FFR in a letter dated March 24, 1989, signed by Robert Q. Cannell, Head of the Department of Agronomy.